

A TOTAL EVIDENCE CLADISTIC ANALYSIS
OF THE HALIOTIDAE
(GASTROPODA: VETIGASTROPODA)

by

Daniel Ludwig Geiger

A Dissertation Presented to the
FACULTY OF THE GRADUATE SCHOOL
UNIVERSITY OF SOUTHERN CALIFORNIA
in Partial Fulfillment of the
Requirement for the Degree
DOCTOR OF PHILOSOPHY
(BIOLOGY)

December 1999

Copyright 1999

Daniel Ludwig Geiger

ACKNOWLEDGMENTS

Entire dissertation

My dissertation committee, composed of co-chairs Jim McLean and Russ Zimmer, and members Dave Bottjer, Bob Lavenberg, and Loren Smith, provided guidance when needed, but otherwise permitted me to pursue my work with minimal interference.

I would like to thank all the staff at museums, who helped to make stays at their institutions successful, helped with library research, and made specimens available for study: Henk Mienis (HUU), Kathie Way, Julia Freeman, Joan Pickering, David Reid, and John Todd (BMNH), Alison Trew (NMW), Yves Finet and Claude Vaucher (MHNG), Roger Pickery (KBIN), Raye Germon, Jerry Harasewych, and Alan Kabat (USNM), Jim McLean and Lindsey Groves (LACM), Paul Scott and Henry Chaney (SBMNH), Terry Gosliner (CASIZ), David Lindberg (UCMP), Paula Mikkelsen (DMNH & AMNH), Gary Rosenberg and Ned Gilmore (ANSP), Peter Jung (NHB), Philippe Bouchet, Pierre Lozouet, and Philippe Maestrati (MNHN), Margaret Gosteli (NMBE), K. Boss (MCZ), Winston Ponder and Ian Loch (AMS), Shirley Slack-Smith (WAM), and Eike Neubert (SMF).

Additional specimens were made available by the following persons: Ted Baer (Lutry, Switzerland), Dominik and Verena Brantschen-Geiger (Bern, Switzerland), Franziska Brantschen (Randa, Switzerland), Anuschka Faucci (Basel, Switzerland), Marc Girona (France), Mark Jones (Auckland, New Zealand), Peter Schuchert (MHNG), Solly Singer (Rehovot, Israel), Brian Hayes (South Africa), Joan Koven (Astrolabe Inc., Washington, D. C.), Roger Pickery (Wilrijk, Belgium), Katharine Stewart (Carmel, California), Buzz Owen (Gualala, California), and Wolfgang Wranik (University of Rostock, Germany).

The systematics discussion group at the Los Angeles County Museum of Natural History was instrumental in shaping my thoughts on cladistics: Bob Bezy, Brian Brown, Kirk Fitzhugh, Guillermo Herrera, John Heynig, David Kizirian, Bob Lavenberg, Don Reynolds, and Christine Thacker.

The staff at Hancock Library at USC and the Research Library of the Natural History Museum of Los Angeles County helped to locate many obscure references.

The work was supported by the following grants and fellowships. The visits of DMNH and ANSP were made possible through a DuPont Merck Scholarship in Malacology, and a Jessup Fellowship, respectively. Research grants from the Hawaiian Malacological Society, the Western Society for Malacology, the Lerner Gray Fund for Marine Research (AMNH), Sigma Xi, and a Trojan League Grant are kindly acknowledged. USC granted a one year Dissertation Fellowship.

This dissertation was written in Word 5.1 and QuarkXPress 4 with Times New Roman as justified font and Courier as monospaced font. Illustrations were made in Photoshop 3, and 4, MacDraw II, Powerpoint 4, QuarkXPress 3.1, and MacClade.

Chapter 1: Recent taxa

This work would not have been possible without the critical remarks by several colleagues, who also made their knowledge and data generously available to me: Phil Colman (Sydney, Australia), Youn-Ho Lee (Pasadena, California), Mark Jones (Auckland, New Zealand), Roger Pickery (Wilrijk, Belgium), Benjamin Singer (Rehovot, Israel), Scoresby A. Shepherd (Adelaide, Australia), Katharine Stewart (Carmel, California), Victor Vacquier (San Diego, California), Rick Fallu (Canberra, Australia), Buzz Owen (Gulala, California), and Luiz Simone (Sao Paulo, Brazil). Jim McLean (Los Angeles, California), Veronica Miller (Los Angeles, California), Scoresby A. Shepherd,

José Leal (Sanibel, Florida), and two anonymous reviewers read the manuscript, including earlier drafts, and made valuable comments. Gert Lindner (Hamburg, Germany) pointed out a typographical error in Geiger (1998a). This chapter has been slightly updated from Geiger (1998a), however, no new taxonomic decisions have been introduced, because a dissertation is unsuitable for such procedures.

Chapter 2: Fossil taxa

This chapter is a collaborative effort by DLG and Lindsey T. Groves (LACM Malacology/Invert. Paleo). LTG has contributed numerous Neogene locality citations, provided obscure references and some figured specimens, and reviewed various ms drafts particularly for paleontologic accuracy. DLG has written and rewritten in excess 90% of the text and produced the figures and plates. This chapter has been published in *Journal of Paleontology* (Geiger & Groves, 1999).

Dave Bottjer (USC), Jim McLean (LACM), and Richard Squires (California State University Northridge), David Lindberg (UCMP), David Dockery, Laurie Anderson, and Tim Hazen critically reviewed various drafts of the manuscript. Akihiro Matsukuma (Kyushu University) kindly helped with some of the obscure references. Dariusz Stramski (USC) and Anders Warén (Swedish Museum of Natural History) translated a Polish and a Swedish paper, respectively, for us. Tita Rodica (Muzeul de Istorie Naturala “Grigore Antipa”, Bucharest) drew our attention to a Romanian publication and translated the relevant parts. Additional information was provided by Claus Hedegaard (UCMP), Hans-Dietrich Laatsch and Manfred Jakubeit (Wolfsburg, Germany), and Padermsak Jarayanbhand (Thailand).

Chapter 3: DNA sequence alignment

This chapter is a collaborative effort between DLG and Kirk Fitzhugh (LACM). The chapter was build on the abductive nature of cladistics worked out by KF. The application of abductive principles to molecular data with the comparison to practices in mophology was carried out by DLG including the development of the new coding strategies. Between 80 and 90% of the text was written by DLG. This chapter is in review with Cladistics since May 1999.

A number of people are thanked for engaging discussions on the above topic. The mention of their names shall not signal their agreement with the content of the chapter. They have been instrumental, however, in sharpening the arguments: Verena and Dominik Brantschen-Geiger, Mark Dawson, Paul Flook, Gonzalo Giribet, Guillermo Herrera, Bob Lavenberg, Jim McLean, Gavin Naylor, Jan Pawlowski, Mason Posner, Mark Siddall, Petra Sierwald, Ellen Strong, and Ward Wheeler. Brian Brown, Doug Eernisse, Jaqueline Reich, and Don Reynolds read earlier versions of the manuscript and made valuable suggestions.

Chapter 4: Biogeography

My appreciation goes to the Palo Alto Research Center of Xerox for offering the free map site on the wold wide web. Jim McLean helped to improve the manuscript. The manuscript is in press with the *Bollettino Malacologico* since August 1999.

Last but not least

The beginnings of this dissertation go back to my highschool years. A number of persons have been instrumental in futhering my interst in marine invertebrates, and particularly in mollusks. Bo Holmstead (Karolinska Institute, Stockholm), Yehuda and

Fanny Mazur (Weizman Institute, Rehovot, Israel), and Prof. emeritus Nüesch (University of Basel) convinced my parents that my interest in snails was worthy of encouragement. And so my family—my mother Anneliese Geiger-Cloos, my father Urs-Peter Geiger, and my sister Verena Brantschen-Geiger—did support in the most admirable fashion the son/brother. Early mentors at the University of Basel include Lukas Hottinger, Peter Jung (NHB), David Senn, and Werner Stingeli. Continuing contact with fellow shell collectors not mentioned elsewhere is a joy as always: Yvonne Grimmer, Roland Hadorn, Jean-Caude Caillez, Michael Trippner, Axel Alf.

Support and friendship with many fellow graduate students were mission critical for this undertaking: Nick Appelman, Jay Vavra, Judy Doino-Lemus, Todd Zimmerman (UCLA), Ellen Strong (GWU), and Rob Guralnick (UCB). Other people responsible for keeping me going include Michel Burla, Janka Jarchow, Jasmin Joshi, Alexander Kocyan, Veronica Miller, Willi Rühl, Willy Rühl, Marlies Rühl, Brigitte Wyss-Heiz.

Table of Contents

INTRODUCTION

CHAPTER 1: RECENT GENERA AND SPECIES OF THE FAMILY HALIOTIDAE RAFINESQUE, 1815 (GASTROPODA: VETIGASTROPODA)

Introduction	3
Materials and methods	4
Systematic affinities and characteristics of the family	6
Problems pertaining to the taxonomy of abalone	8
Tremata as a taxonomic character and teratological type specimens	8
Hybrids	12
Genus-level taxa	16
<i>Haliotis, sensu stricto</i>	16
Other genus-level taxa, and their type species	18
Use of genus-level taxa	21
Species-level taxa	22
Use of subspecies	22
Index	25
Notes	39
Caribbean	69
European and Senegalese	70
South African	70
Eastern African, Red Sea and Persian Gulf	71

Indian Ocean and Tropical West Pacific	72
Temperate Australian	73
New Zealand	74
Northwestern Pacific	75
Northeastern Pacific	75
Tropical Eastern Pacific	76
Zoogeography	76

**CHAPTER 2: REVIEW OF FOSSIL ABALONE
(GASTROPODA: VETIGASTROPODA: HALIOTIDAE)
WITH COMPARISON TO RECENT SPECIES.**

Introduction78
Diagnostic Characters of the Family78
Taxonomy80
The shell as the basis of taxonomy	80
Fossil abalone taxa	83
Fossil abalone in the phylogenetic context	88
Preservation89
Paleoenvironments90
Reef paleoenvironments	90
Shallow-water rocky paleoenvironments	91
Time Range92
Geographic Distribution94

**CHAPTER3: DNA SEQUENCE ALIGNMENT IN THE LIGHT OF THE
ABDUCTIVE NATURE OF CLADISTIC HYPOTHESIS**

GENERATION.

Introduction	.111
Molecular and Morphological Character Equivalence	.112
The observational phase	113
The explanatory phase	116
Cladistics as science	119
Observation: From Perception to Classification	.121
i: Special similarity	123
ii: Positional correspondence	123
Sequence alignment	124
Observation and Explanation	.125
Primary and secondary homology	125
Separation of power	126
Character Independence	.128
Objectivity and Subjectivity	.131
Character selection and observation	131
Alignment parameters	132
Manual editing	133
Alternative Coding Strategies	.133
Elision	134
Case sensitive	136
Missing data	137

Polymorphic	137
Exclusion	138
Contraction	138
Homology is Special Similarity <i>sensu</i> Remane (1952)	139
Belief formation	140
Rational for data exclusion	142
Highly Dissimilar Taxa	146
Stretch coding	148
Block coding	149
Presence/absence coding	150
Comparison to practice in morphology	150
Better alignment?	152
Conclusions and Recommendations	154

**CHAPTER 4: DISTRIBUTION AND BIOGEOGRAPHY OF THE
RECENT HALIOTIDAE (GASTROPODA: VETIGASTROPODA)
WORLD-WIDE.**

Introduction	156
Materials and Methods	158
Sources of Raw Data	158
Abbreviations of collections	159
Biogeographical regions	160
Biogeographical analysis	162
Cladistic analysis of taxa	165

Distribution	174
Distributional patterns	175
Taxonomic issues	177
Area cladograms	178
Cladistic analysis of taxa	182
Illustrations of species.	185
Biogeography and the fossil record	188
DISTRIBUTIONAL DATA AND MAPS	189
Mediterranean and West African Species (Figures 4-4-14, 4-146, 4-147, 4-149, 4-150, 4-170, 4-172, 4-174)	189
<i>Haliotis marmorata</i> Linnaeus, 1758 (Figures 4-14, 4-146, 4-149)	189
<i>Haliotis stomatiaeformis</i> Reeve, 1846 (Figures 4-14, 4-147, 4-150)	192
<i>Haliotis t. tuberculata</i> Linnaeus, 1758 (Figures 4-4-9, 4-12-14, 4-172, 4-174) .	192
South African Species (Figures 4-14, 4-142-145, 4-148, 4-151-156, 4-159-162)	200
<i>Haliotis midae</i> Linnaeus, 1758 (Figures 4-14, 4-142-145, 4-148)	200
<i>Haliotis parva</i> Linnaeus, 1758 (Figures 4-14, 4-154-156)	201
<i>Haliotis queketti</i> Smith, 1910 (Figures 4-14, 4-161, 4-162)	202
<i>Haliotis spadicea</i> Donovan, 1808 (Figures 4-14, 4-151-153)	202
<i>Haliotis speciosa</i> Reeve, 1846 (Figures 4-14, 4-159, 4-160)	203
East African Species (Figures 4-14-23, 4-130, 4-133, 4-157, 4-158, 4-169, 4-171)	
204	
<i>Haliotis mariae</i> Gray, 1826 (Figures 4-14-16)	204
<i>Haliotis pustulata</i> Reeve, 1846 (Figures 4-14, 4-17-20)	204
<i>Haliotis rugosa</i> Lamarck, 1822 (Figures 4-14, 4-21, 4-22)	208
<i>Haliotis squamosa</i> Gray, 1826 (Figures 4-14, 4-130, 4-133)	208

<i>Haliotis unilateralis</i> Lamarck, 1822 (Figures 4-14, 4-23, 4-157, 4-158, 4-169, 4-171)	208
Indo-Pacific and Central Pacific Species (Figures 4-24-44, 4-63, 4-66, 4-116, 4-117, 4-119, 4-120, 4-122-129, 4-136-141, 4-163-166, 4-175, 4-176)	209
<i>Haliotis asinina</i> Linnaeus, 1758 (Figures 4-35, 4-36, 4-43, 4-165, 4-166)	210
<i>Haliotis clathrata</i> Reeve, 1846 (<i>non</i> Lichtenstein, 1794) (Figures 4-24, 4-138, 4-141, 4-175, 4-176)	219
<i>Haliotis crebrisculpta</i> Sowerby, 1914 (Figures 4-24, 4-25, 4-28)	222
<i>Haliotis diversicolor</i> Reeve, 1846 (Figures 4-24, 4-40-42)	222
<i>Haliotis dissona</i> (Iredale, 1929) (Figures 4-24, 4-139, 4-140)	224
<i>Haliotis dohrniana</i> Dunker, 1863 (Figures 4-24, 4-120, 4-123)	225
<i>Haliotis fatui</i> Geiger, 1999 (Figures 4-43, 4-116, 4-117)	225
<i>Haliotis glabra</i> Gmelin, 1791 (Figures 4-26, 4-27, 4-29, 4-30, 4-43)	227
<i>Haliotis jacnensis</i> Reeve, 1846 (Figures 4-24, 4-124-129)	228
<i>Haliotis ovina</i> Gmelin, 1791 (Figures 4-31-33, 4-43)	230
<i>Haliotis planata</i> Sowerby, 1882 (Figures 4-43, 4-119, 4-122)	236
<i>Haliotis pulcherrima</i> Gmelin, 1791 (Figures 4-38, 4-39, 4-44)	237
<i>Haliotis rubiginosa</i> Reeve, 1846 (Figures 4-24, 4-136, 4-137)	239
<i>Haliotis squamata</i> Reeve, 1846 (Figures 4-24, 4-63, 4-66)	240
<i>Haliotis varia</i> Linnaeus, 1758 (Figures 4-34, 4-37, 4-43, 4-163, 4-164)	241
Australian Endemic Species (Figures 4-45-67, 4-131, 4-132, 4-134, 4-135, 4-183-192)	250
<i>Haliotis brazieri</i> Angas, 1869 (Figures 4-45, 4-48, 4-67)	250
<i>Haliotis coccoradiata</i> Reeve, 1846 (Figures 4-47, 4-50, 4-67, 4-186)	253
<i>Haliotis cyclobates</i> Péron & Lesueur, 1816 (Figures 4-51, 4-52, 4-67, 4-185)	254

<i>Haliotis elegans</i> Philippi, 1844 (Figures 4-53, 4-54, 4-67, 4-192)	255
<i>Haliotis hargravesi</i> Cox, 1869 (Figures 4-46, 4-49, 4-67, 4-189)	256
<i>Haliotis laevigata</i> Donovan, 1808 (Figures 4-57, 4-60, 4-67, 4-184)	256
<i>Haliotis roei</i> Gray, 1826 (Figures 4-61, 4-64, 4-67, 4-188)	257
<i>Haliotis rubra</i> Leach, 1814 (Figures 4-62, 4-65, 4-67, 4-183)	260
<i>Haliotis scalaris</i> Leach, 1814 (Figures 4-55, 4-56, 4-58, 4-59, 4-67, 4-190) . . .	263
<i>Haliotis semiplicata</i> Menke, 1843 (Figures 4-67, 4-131, 4-132, 4-134, 4-135, 4-187)	264
New Zealand Species (Figures 4-67-79, 4-177-182)	266
<i>Haliotis australis</i> Gmelin, 1791(Figures 4-67, 4-68, 4-71, 4-177, 4-178)	266
<i>Haliotis iris</i> Gmelin, 1791 (Figures 4-67, 4-74-79, 4-179, 4-180)	268
<i>Haliotis virginea</i> Gmelin, 1791 (Figures 4-67, 4-69, 4-70, 4-72, 4-73, 4-181, 4-182)	270
Northwest Pacific Species (Figures 4-80-86, 4-118, 4-121)	272
<i>Haliotis discus</i> Reeve, 1846 (Figures 4-80, 4-83, 4-86)	272
<i>Haliotis exigua</i> Dunker, 1877 (Figures 4-87, 4-118, 4-121)	276
<i>Haliotis gigantea</i> Gmelin, 1791 (Figures 4-81, 4-84, 86)	276
<i>Haliotis madaka</i> Habe, 1977 (Figures 4-82, 4-85, 4-86)	277
East Pacific Species (Figures 4-87-102, 4-104-107, 4-109, 4-110, 4-112, 4-113, 4-167, 4-168, 4-173)	278
<i>Haliotis corrugata</i> Wood, 1828 (Figures 4-88, 4-91, 4-100)	278
<i>Haliotis cracherodii</i> Leach, 1814 (Figures 4-89, 4-92, 4-100)	284
<i>Haliotis dalli</i> Henderson, 1915 (Figures 4-107, 4-110, 4-113)	289
<i>Haliotis fulgens</i> Philippi, 1845 (Figures 4-90, 93, 100, 173)	290
<i>Haliotis k. kamtschatkana</i> Jonas, 1845 (Figures 4-87, 94, 97)	293

<i>Haliotis kamtschatkana assimilis</i> Dall, 1878 (Figures 4-95, 4-99, 4-100)	296
<i>Haliotis roberti</i> McLean, 1970 (Figures 4-107, 4-109, 4-112, 4-167, 4-168) . .	298
<i>Haliotis rufescens</i> Swainson, 1822 (Figures 4-96, 4-99, 4-100)	298
<i>Haliotis sorenseni</i> Bartsch, 1940 (Figures 4-100, 4-101, 4-104)	303
<i>Haliotis walallensis</i> Stearns, 1898 (Figures 4-100, 4-102, 4-105)	305
Western Atlantic Species (Figures 4-107, 4-108, 4-111, 4-114, 4-115)	307
<i>Haliotis aurantium</i> Simone, 1998 (Figures 4-107, 4-108, 4-111)	307
<i>Haliotis pourtalesii</i> Dall, 1881 (Figures 4-107, 4-114, 4-115)	307

**CHAPTER 5: A TOTAL EVIDENCE CLADISTIC ANALYSIS OF THE
WORLD-WIDE HALIOTIDAE.**

Introduction	324
Materials and Methods	325
Source of specimens	325
SEM preparation	325
Cladistic methodology	325
Outgroup comparison	329
Allozyme frequencies	330
Outgroup comparison	332
Reanalysis of allozyme data with comparison to the results of Brown (1993) .	332
DNA coding strategies for 16kD cDNA	336
Outgroup comparison	337
Reanalysis of the lysin data	337

Partial 16S mtDNA (Charley Wray, unpubl. data)	.345
Morphological characters	.348
The radula in abalone	348
Outgroup comparison	351
Characters and character states	352
The Epipodium	.356
Outgroup comparison	359
Characters and character states	359
Potential characters not used	363
Shell morphology	363
Shell mineralogy	363
Analyses with morphological characters	.364
Classification	.369
<i>Haliotis, sensus stricto</i>	370
<i>Nordotis</i>	370
<i>Notohaliotis</i>	371
<i>Sanhaliotis</i>	372
Use of Genus-level taxa	372
CONCLUSIONS	
Chapter 1: Recent Taxa	.373
Chapter 2: Fossil Taxa	.373
Chapter 3: DNA Sequence Alignment	.373
Chapter 4: Biogeography	.374
Chapter 5: Total Evidence Cladistic Analysis	.375

List of Figures

Figures 1-2 - 1-5. Shells of imperforate specimen and potential hybrid of <i>Haliotis bra-</i> <i>zieri</i> x <i>hargravesi</i>	11
Figures 1-6 - 1-9. Shells of designated lectotypes for <i>Haliotis multiperforata</i> and <i>H.</i> <i>revelata</i>	29
Figures 1-10 - 1-15. Shells of uncommonly illustrated abalone I.	53
Figures 1-16 - 1-21. Shells of uncommonly illustrated abalone II.	54
Figures 1-22 - 1-27. Shells of uncommonly illustrated abalone III.	55
Figures 1-28 - 1-33. Shells of uncommonly illustrated abalone IV.	56
Figure 2-1: Specimens of fossil haliotids	79
Figure 2-2: Histograms of number of open tremata for several species of <i>Haliotis</i>	82
Figure 2-3. Localities of fossil abalone listed in Appendix.	95
Figure 3-1. Treatment of morphological and molecular characters in cladistics with respect to homology.	114
Figure 3-2. The effect of character coding on homology statements in questionably aligned DNA sequences, using hypothetical sequences of four taxa.	135
Figure 3-3. Coding strategies for a few highly dissimilar taxa.	147
Figure 4-1. Illustration of the three biogeographical models proposed for the origin of <i>Haliotidae</i>	157
Figure 4-2. Area cladogram (Brooks parsimony) for the genus <i>Haliotis</i>	179
Figure 4-3. Midpoint rooted taxon cladogram of the genus <i>Haliotis</i>	183
Figure 4-4-13. Shells of the European-Mediterranean species of <i>Haliotis</i> spp.	190
Figure 4-14. Distribution of European-Mediterranean, west African, and south African species of <i>Haliotis</i> spp.	191
Figures 4-15-23. Shells of East African abalone.	205

Figure 4-24. Distribution of Indo-Pacific species of <i>Haliotis</i> sp. (I).	211
Figures 4-25-42. Shells of Indo-Pacific abalone.	212
Figure 4-43. Distribution of Indo-Pacific species of <i>Haliotis</i> spp. (II).	226
Figure 4-44. Distribution of the central Pacific (Tuamotus) <i>Haliotis pulcherrima</i>	238
Figures 4-45-60. Shells of abalone endemic to Australia I.	251
Figures 4-61-66. Shells of abalone endemic to Australia II.	258
Figure 4-8. Distribution of endemic Australian and New Zealand species of <i>Haliotis</i> spp.	265
Figures 4-68-79. Shells of New Zealand abalone.	267
Figures 4-80-85. Shells of the northwest Pacific abalone.	273
Figure 4-86. Distribution of northwest Pacific species of <i>Haliotis</i> spp.	274
Figure 4-87. Distribution of east Pacific <i>Haliotis k. kamtschatkana</i>	279
Figures 4-88-99. Shells of the northeast Pacific abalone I.	280
Figure 4-100. Distribution of east Pacific species of <i>Haliotis</i> spp. (I).	299
Figures 4-101-106. Shells of Northeastern Pacific abalone I, and unidentified species. 304	
Figure 4-107: Distribution of east Pacific (II), tropical Pacific, and western Atlantic species of <i>Haliotis</i> spp.	308
Figure 5-1. Trees generated from allozyme frequency data (Brown, 1993).	333
Figure 5-2. Trees generated from data of Lee & Vacquier (1995).	338
Figure 5-3. Strict consensus topology of 16S mtDNA sequences.	346
Figure 5-4. Strict consensus tree from all lysin, allozyme, and 16S mtDNA data com- bined.	347
Figure 5-5. Diagram illustrating the terminology used for the radular teeth found in the family Haliotidae.	350

Figure 5-6. Illustration of radular character states for central field. Number on images refer to character number.	353
Figure 5-7. Illustration of radular character states for lateral teeth 3-5.	354
Figure 5-8. Illustration of radular character states for marginal teeth.	355
Figure 5-9. Sample epipodia of <i>Haliotis</i> spp.	358
Figure 5-10. Example of a hypobranchial gland.	362
Figure 5-11. Strict consensus tree of 680 MPRs from combined analysis of allozyme, lysin, 16S mtDNA, and morphology.	365
Figure 5-12: Strict consensus tree of 32 ingroup taxa from combined analysis of allozyme, lysin, 16S mtDNA, and morphology, without outgroups.	367
Figure 5-13. Strict consensus tree for combined analysis of lysin, 16S mtDNA, and morphology, without allozyme data.	368

List of Tables

Table 1-1. Genus-level taxa in the family Haliotidae.	17
Table 2. Differences between the valid taxa of this study with their status in Wagner & Abbott (1978) and illustrations in Kaicher (1981).	26
Table 2-1: Comparison of upper Tertiary species with Pleistocene-Recent species from California.	85
Table 2-2: Chromosome number in Recent <i>Haliotis</i> spp. as indicated in the respective source.	97
Table 2-3: List of fossil abalone taxa.	100
Table 4-1. Data matrix of biogeographical analysis.	163
Table 4-2. Recoded allozyme frequency data from Brown (1993).	166
Table 4-3. Recoded sequence of the lysin data of Lee & Vacquier (1995).	168
Table 5-1: Source of specimens and data.	326
Table 5-2. Character states for the radular characters.	349
Table 5-3: Character states for epipodial characters.	357
Table 5-4. Character states for hypobranchial gland characters.	361